Chapter 10 / Capítulo 10

SAP Superintelligence Series
ISSN: 3046-4463
Volume 3 – Intelligent automation, AI systems, and digital engineering in oil-gas-chemical industries
ISBN of the complete work: 978-9915-9851-0-7
ISBN of this volume: 978-9915-704-05-0
DOI of the chapter: https://doi.org/10.62486/978-9915-9851-0-7_202641
Pages: 167-183

©2026 The authors. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) 4.0 License.

ORIGINAL

Engineering Transformation at the Industrial-Quantum Interface: Opportunities for Topological Superconductive Materials in Oil, Gas, and Chemical Systems

Transformación de la ingeniería en la interfaz industrial-cuántica: oportunidades para materiales superconductores topológicos en sistemas de petróleo, gas y químicos

S. Dhamotharan[1] ✉, M. Muthukrishnan[2] ✉, M. Bharath[1] ✉, P. Mathankumar[1] ✉, N. Ragavan[1] ✉, P. Prabhu[1] ✉, Shafi Danyalov[2] ✉

[1]Department of Mechanical Engineering, KIT-Kalaignarkarunanidhi Institute of Technology, Kannampalayam. Coimbatore, Tamil Nadu, 641 402, India.

[2]Department of Mechanical Engineering, Christ the King Engineering College, Karamadai. Coimbatore, Tamil Nadu, 641 104, India.

[3]Azerbaijan University of Architecture and Construction. Baku, Azerbaijan.

Corresponding Author: S. Dhamotharan ✉

ABSTRACT

The oil, gas, and chemical sectors are experiencing a fast technological change that is mainly influenced by the increasing operational complexity, the strict environmental requirements, and the rising demand for high-precision, data-centric engineering solutions. At the same time, significant advances in condensed-matter physics, in particular, the identification and characterization of topological superconductive materials, are leading to quantum technologies that, in the future, may have extraordinary computational power, sensing accuracy, and security. Topological superconductors, which essentially incorporate superconductivity with nontrivial topological order to yield exotic quasiparticles like Majorana fermions, are among the most intriguing quantum materials to be considered. Hence, these materials that interconnect quantum computation with condensed matter physics, form a quantum information processing platform that is compatible with fault-tolerance. If one assumes spin–orbit coupling, time-reversal symmetry, and unconventional pairing as the basic ideas, topological superconductivity can be discovered in an extremely wide variety of systems such can one as heavy-fermion metals, non-centrosymmetric compounds, engineered heterostructures, and topological insulators. The comprehension of topological phases and protected surface states has also been facilitated by theoretical contributions of Kane, Qi, Zhang, and other researchers. Meanwhile, experimental work on materials such as Cu-doped Bi_2Se_3, Sr_2RuO_4, and UTe_2 has produced strong evidence of unconventional pairing and symmetry breaking. The versatility of topological superconductivity is further demonstrated by ongoing studies in Dirac and Weyl semimetals, superconducting carbides, and hybrid nanowires.This paper examines the theoretical foundations, material realizations, and potential quantum applications of topological superconductors.

Keywords: Topological Superconductors; Majorana Fermions; Quantum Computation; Spin–Orbit Coupling; Unconventional Pairing; Topological Order.

RESUMEN

Los sectores del petróleo, el gas y la química están experimentando una rápida transformación tecnológica, influenciada principalmente por la creciente complejidad operativa, los estrictos requisitos ambientales y la creciente demanda de soluciones de ingeniería de alta precisión y centradas en datos. Simultáneamente, los avances significativos en la física de la materia condensada, en particular la identificación y caracterización de materiales superconductores topológicos, están dando lugar a tecnologías cuánticas que, en el futuro, podrían ofrecer una potencia computacional, una precisión de detección y una seguridad extraordinarias. Los superconductores topológicos, que esencialmente incorporan superconductividad con un orden topológico no trivial para producir cuasipartículas exóticas como los fermiones de Majorana, se encuentran entre los materiales cuánticos más interesantes a considerar. Por lo tanto, estos materiales, que interconectan la computación cuántica con la física de la materia condensada, conforman una plataforma de procesamiento de información cuántica compatible con la tolerancia a fallos. Si se asume el acoplamiento espín-órbita, la simetría de inversión temporal y el emparejamiento no convencional como ideas básicas, la superconductividad topológica puede descubrirse en una amplia variedad de sistemas, como metales con fermiones pesados, compuestos no centrosimétricos, heteroestructuras diseñadas y aislantes topológicos. La comprensión de las fases topológicas y los estados superficiales protegidos también se ha visto facilitada por las contribuciones teóricas de Kane, Qi, Zhang y otros investigadores. Mientras tanto, el trabajo experimental en materiales como Bi_2Se_3 dopado con Cu, Sr_2RuO_4 y UTe_2 ha producido evidencia sólida de emparejamiento no convencional y ruptura de simetría. La versatilidad de la superconductividad topológica se demuestra aún más mediante estudios en curso sobre semimetales de Dirac y Weyl, carburos superconductores y nanocables híbridos. Este artículo examina los fundamentos teóricos, las realizaciones materiales y las posibles aplicaciones cuánticas de los superconductores topológicos.

Palabras clave: Superconductores Topológicos; Fermiones de Majorana; Computación Cuántica; Acoplamiento Espín-Órbita; Emparejamiento no Convencional; Orden Topológico.

INTRODUCTION

The usual class of crystalline solids, magnets, and superconductors with specific unique states fall within the category of quantum matter. By looking at their spontaneous symmetry breaking, these states can be categorized.[1] By disrupting this uniformity or reducing the number of points of view invariance, symmetry breaking creates a more organised and unlikely state.[2] Topologyin math is the study of the shape and space. The characteristics of the space that are still the same, if it is changed by the continuous transformations, like elongation, rotation and folding, are what the work is about.[3] Topology is a concept in mathematics used to group different shapes. The area of mathematics known as topology examines the characteristics of shapes that are unaffected by deformations; in other words, a geometrical object maintains its shape despite ongoing deformations. Sphere and 1-torus are well-known examples of objects having non-trivial topology.[4] It focuses on properties that do not change even when the shape is distorted and is an important part of modern mathematics. Topology has applications in fields such as physics, chemistry, economics, and computer science.[5] Unusual superconductors with a bulk superconducting gap and in-gap Majorana states on the border are known as topological superconductors (TSCs). For quantum processing, they could be used as topological qubits.[6] Superconductivity is often explained by the Ginzburg-Landau(G-L) hypothesis. The free energy, F, of a superconductor close to the superconducting transition can be expressed

using the complex order parameter field, −r =|ψr|eiφ(r). Similar to a wave function in quantum physics, the quantity |ψr2| is a measure of the local density, and below a phase transition into a superconducting state, ψr is nonzero.[7] The free energy is shown as, assuming smallness of and smallness of gradients:

$$F = F_n + \alpha|\psi|^2 + \frac{\beta}{2}|\psi|^4 + \frac{1}{2m^*}|(-i\hbar\nabla - e^*\mathbf{A})\,\psi|^2 + \frac{|\mathbf{B}|^2}{2\mu_0}$$

Where Fn is the normal phase free energy, α and ß are phenomenological parameters, m* is an electron's effective mass, e* is its effective charge, A is the magnetic vector potential, and B is the magnetic field. The Ginzburg-Landau equations are derived by minimizing the free energy with respect to variations in the vector potential and order parameter. Ginzburg Landau (G-L) theory may be used to identify this order parameters and provides a distinctive description of these quantum states. However, since a quantum material with an order parameter creates a new quantum state, topology transcends the conceptual framework of Ginzburg Landau Theory.[8] In the 1980s, Von Klitzing discovered Quantum Hall (QH) states, a class of quantum states in the 2-D system. This system features an insulating gap in the center and a conducting edge due to the skipping orbits produced by the externally provided magnetic field.[9] These edge states provide an electric current with a unidirectional, dissipation-free channel, which results in the Quantized Hall Effect (QHE). Topological invariants are characteristics of topological phases that do not affect the bulk energy gap in the surface states and are resistant to smooth deformation. The unusual topological phases, which are also strongly influenced by the energy gap in bulk, result in the creation of two new classes of quantum materials: Topological Insulators (TIs) and Topological Superconductors (TSCs).[10,11] Topological superconductivity is related with Majorana fermions (MFs), which are quasiparticle excitations. In 1937, EttoreMajorana speculated that there could be a particle that is its own antiparticle.[12] In the quantum Hall effect, which is observed in complex electron systems exposed to high magnetic fields and low temperatures, Thouless, Kohmoto, den Nijs, and Nightingale developed a groundbreaking connection between topological invariants and Hall conductivity.[13] The success of topological invariants in bulk quantum solid state systems is followed by ongoing research on the Hall Effect and superfluid. Topological invariants are topological space quantities that have discrete values (typically normalised to be integer-valued) and do not change under continuous deformations of the space. TheMajorana fermion is a hypothesised collective electronic state that behaves like an antiparticle particle.[14,15] Since Thouless, Kosterlitz, and Haldane received the 2016 Nobel Prize in Physics for their groundbreaking discoveries in topological quantum matter and topological phase transitions, the experimental signature of these peculiar phases of matter has been established.[16] Since an MF is half of a normal fermion, a superposition of two geographically separated MFs yields a fermionic state.[17] This is usually only a mathematical operation with no physical implications because the two MFs are physically situated close to one another, greatly overlap, and cannot be handled independently.Such highly delocalized fermionic states are resistant to local perturbations that impact only one of the Majorana components and are resistant to the majority of kinds of decoherence.[18] The non-Nabelian statistics of MF, on the other hand, imply that states can be transformed by physical exchanges, giving rise to the notion of low-decoherence topological quantum computing.[19] These industries (oil, gas, and chemical) are moving into a time where the usual measurement and modeling tools cannot manage the complexities of extraction, refining, and environmental compliance that are growing. The use of advanced materials and nanoscale engineering is changing the industrial diagnostics area; to illustrate, better nanoparticle synthesis and characterization have allowed structural analysis

to be more accurate and predicting material performance in extreme conditions, as can be seen in the studies that have been done recently on size-dependent XRD signatures in monodisperse metal and oxide nanostructures.[20] These changes highlight the increasing need for very detailed measurements to be able to follow the process, evaluate the catalyst, and locate corrosion—in other words, to increase slightly the measurement fidelity to have a significant result in operational efficiency. At the same time, the quantum technology development is another fundamentally different way to resolve the long-standing problems of sensing and computation in energy systems. In particular, quantum-enhanced sensors have demonstrated the capability to pick up very weak electromagnetic, thermal, and gas-phase signals with a sensitivity level that is many times better than that of classical instruments. Hence, the door is open to new, timely, and cost-effective ways of methane leak detection, structural-health monitoring of pipelines, and early anomaly identification in refinery units.[21,22] Quantum algorithms and hybrid quantum-classical simulations are also considered as means to speed up subsurface modelling, materials discovery, and process optimization in the energy sector.[23] The integration of these technologies is additionally supported by the impetus of recent extensive assessments of methane emissions that underline the necessity for accurate and continuous monitoring systems as a way to fulfill regulatory and environmental commitments.[24] In combination, these changes indicate a move to an industrial–quantum convergence where nanoscale characterization, quantum sensing, and quantum simulation might, by their interdependence, revolutionize the technological bases of the oil, gas, and chemical industries.

METHOD

Majorana Fermions

Majorana fermions are topologically shielded, making them resistant to environmental thermal or electrical disturbances. Majorana fermions promise zero-resistance information and communications technology, addressing contemporary electronics' contributing to more than 8 % global energy usage and providing a sustainable future for computing.[5] Moreover, topological superconductors are the main potential materials for achieving topological quantum computing due to the existence of Majorana zero-energy modes.[19,22] The scientific community is interested in superconductivity in topological materials because of the presence of Majorana fermions, which are conceptualized from quantized thermal conductivity, a zero-biased conduction peak, and the anomalous Josephson Effect.[21]




Figure 1. Schematic Illustration of Proximity-Induced Superconductivity in a Spin-Orbit Coupled Material

S-wave topology as shown in figure 1 has one conducting channel, whereas P-wave topology has two conducting channels. D-wave topology is characterised by four conducting channels.[15] The quantity of conducting channels and the electric current's direction are the primary distinctions. S-wave and P-wave topologies are employed for signal transmission, whereas D-wave topology is utilised for power transmission.

A superconducting metal and a semiconductor are frequently combined to form topological superconductors and Majorana fermions, are produced through interactions between these materials. Topological materials have gotten a lot of scholarly interest during the last few decades. High-efficiency thermoelectrics, current-induced switching for spintronic applications, energy-free electronic states like the quantum spin Hall effect and the quantum anomalous Hall effect, topological dipoles for future photovoltaic and photodetectors, and other energy conversion and storage techniques have all been shown at the lab scale.

TSC was classified as either weak or powerful. Superconductors with a superconducting gap and topological features were considered weak TSCs. A complete superconducting gap with odd parity pairing symmetry, non-conducting gapless surface states, and superconducting vortex cores with Majorana Zero Modes (MZMs) were the three distinctive characteristics of the strong TSC.

TSC can also be classified into two groups based on time-reversal symmetry: those that preserve TRS and those that break it. Detailed classification can also be done based on whether TRS is broken or preserved. A magnetic field inside superconductors with odd parity pairing is involved in the breakdown of TRS in the former kind of TSC. Due to strong SOC, the TRS in the subsequent class is preserved.[20]




Figure 2. Schematic of a Hybrid Nanowire–Superconductor Device for Majorana Zero Mode Detection

In figure 2 Nanowire (yellow/green tube structure): this is a semiconducting nanowire, often made from materials like InSb or InAs. It is a crucial platform for realizing topological superconductivity.

Superconductor (green block): a superconducting material (commonly Al or NbTiN) is deposited on part of the nanowire. Through the proximity effect, it causes the nanowire to become superconductive.

Normal metal contacts (N, orange blocks): these are metallic leads used to inject and detect electrons from the nanowire.

Gate electrode (grey base labeled “Gate”): used to control the chemical potential in the nanowire, tuning its electronic properties.

Magnetic field (blue arrow labeled $\vec{B}$): applied perpendicular or parallel to the wire to induce a topological phase transition necessary for the emergence of Majorana zero modes.

Majorana bound states (yellow spikes): these are localized zero-energy states at the ends of the superconducting segment, characteristic of a topological superconductor. The symmetries and topological characteristics of the system are used to categorize gapped topological phases of matter. These classes have been studied in a variety of contexts, from two-dimensional systems

to the recently discovered three-dimensional systems. The classification of gapped topological phases and their corresponding topological invariants can be used to determine the nature of the phase transition between different topological states. The symmetries and topological characteristics of the system are used to categorize gapped topological phases of matter.[21]




Figure 3. Electronic Band Structure and Fermi Surface Signatures in Superconductors

In figure 3 Green Cone: labeled as the "Light" cone, representing the boundary beyond which quasiparticles cannot escape due to relativistic constraints. Blue Paraboloid (Bulk band): Represents the bulk electronic band structure of the material. It lies within the light cone, signifying allowable electronic states.

TSC topological types and classifications of gapped topological phases of matter




Figure 4. Key Ingredients and Pathways Toward Intrinsic Topological Superconductors

First, a symmetry constraint can be used to categorize the (gapped) topological phases. The kind of topological phase is determined by the symmetry of the Hamiltonian, which is connected to the symmetry constraint. Additionally, there are other kinds of topological phases, including topological crystalline insulators, topological superconductors, and topological insulators. The time-reversal symmetry of the Hamiltonian gives rise to a distinct set of restrictions for the TSC topological classes as shown in figure 4. Second, even-parity and odd-parity topological classes of topological superconductors (TSCs) can be distinguished.[13]

Materials for Topological Superconductors

The realization of TSC experimentally is only limited to a few candidate materials. The difficulty is due to the materials having to be low-temperature superconductors and topological insulators, a combination that is both rare and challenging to obtain. It has been quite a challenge to find the right combination of materials. Researchers have been experimenting with different materials to produce this special combination, but as yet there has been no success. Scientists are now looking for newly discovered materials that can satisfy the requirements. TSCs can be either natural or artificial. Natural TSCs may theoretically be the first TSCs to be exploited for the development of TSC systems, as they offer a safe, inexpensive environment for research and development.[22] Furthermore, they may also serve as a source of providing a more localized approach to TSC deployment and facilitating the creation of custom TSC solutions for specific purposes. However, in order to produce artificial TSCs, 2D heterostructures or 1D wires with several constituent components would need to be synthesized at low temperatures and with great crystallinity.




Figure 5. Electronic Structure and Quantum Well States in Pb/TiBiSe2 and Pb/Si Heterostructures

This is because these structures must be constructed layer by layer or by carefully regulating the nanostructures' composition and geometry. Low temperature synthesis techniques like chemical vapor deposition or molecular beam epitaxy are the only ways to accomplish this.The capacity of these materials to self-assemble and create intricate structures is the reason behind this. They are also easily made in big quantities and are reasonably priced.[10]

In figure 5 (a) crystal structure model of the Pb/$TlBiSe_2$ heterostructure along the (111) direction, showing atomic layers of Pb, Tl, Bi, and Se. (b) Schematic of the surface Brillouin zone (BZ) alignment between Pb (green) and $TlBiSe_2$ (red). (c–d) Measurements of bare $TlBiSe_2$ using angle-resolved photoemission spectroscopy (ARPES): (c) The intensity surrounding the Γ point on the Fermi surface map in momentum space. (d) Band dispersion along the Γ–M direction showing topological surface states (SS), Dirac point (DP), and conduction band (CB). (e–f) ARPES results for Pb deposited on $TlBiSe_2$: (e) Fermi surface of Pb/$TlBiSe_2$. (f) Quasiparticle well states (QWS) and subband formation in the Pb film due to confinement. (g–h) ARPES data for Pb grown on Si substrate: (g) Fermi surface of Pb/Si. (h) Clear QWS subband structure similar to Pb/$TlBiSe_2$, indicating confined electron behavior. Insets: Circular Fermi surface and schematic band dispersions are shown to provide further context for electronic states.vColor scale: Spectral intensity (yellow: high, red: low) representing electron occupancy.

Artificial Candidates

Temperature-stable capacitors (TSCs) are one of the energy storage devices, which can work under extremely high or low temperatures. Besides being made to last longer and be more energydense than regular capacitors, they also release more system reliability. Hence, the first TSCs have been widely engaged as power units for e-mobility and green energy.

The hybridization of a superconductor and a semiconductor by coupling magnetic atoms with a semiconductor nanowire to create a 1D topological superconductor (TSC) is the main result of this research. The mix of these materials is crucial because it allows electrons to flow freely between them and ultimately results in the creation of the superconducting state. A topological superconductor is a minimal energy system comprising a semiconductor nanowire and magnetic atoms that either form a chain or a superconductor. Because the nanowire can move electrons considerably more quickly than conventional materials, scientists have shown that it may be a less energy-consuming form of superconductivity than the present systems. To achieve the best of both worlds, such a system hybridizes a semiconductor nanowire with a magnetic atom chain or a superconductor. The nanowire is the one that carries the current, while the magnetic atoms chain or superconductor provides the necessary magnetic properties for current control. The different charge states can be switched through this device, which in turn allows energy transfers to be more efficient.

This system's indium arsenide (InAs) nanowire shows Zeeman splitting and significant spin-orbit coupling (SOC), making it a prime candidate for a hybrid superconductor (SC) made up of a chain of magnetic atoms and a semiconductor nanowire.Aluminum superconductor in the proximity of the nanowire is, therefore, a relief from dissipation.[8] It turns out that MZMs are anticipated to emerge at the two ends of the wire in such a system.

The MZMs that result from the proximity-induced superconductivity in the (InAs) nanowire are responsible for this operation. When a little magnetic field is introduced along the wire, the Zeeman splitting of the nanowire causes the MZMs to have different energies. Therefore, the zero-bias conductance peak is an indication that the MZMs are tunneling from one end of the wire to the other.[9]

Figure 6 scanning tunneling microscopy/spectroscopy (STM/STS) utilizes a superconducting niobium (Nb) tip to locally measure the UTe_2 sample. Various steps of the experiment are illustrated in the figure. Initially, in the left panel, the Nb tip is shown moving towards the UTe_2 surface, and the tunneling process is depicted. The central panel represents spatial modulation

of the Cooper pair density in UTe_2. Finally, the right panel demonstrates a modulation in the CD. The relative phase between the pair density wave and the CDW is $\Delta\phi = \pi$, which shows that these two are in an antiphase relation.[10]




Figure 6. Visualization of Electron Pair and Charge Density Waves in UTe_2 Using a Niobium Tip

2D TSC

Fu and Kane forecasted the creation of a two-dimensional topological superconductor from an "s-wave superconductor-topological insulator (TSC) heterostructure". Their theoretical work led to a revival of interest in 2D topological superconductors and further research activities followed.




Figure 7. Probing Superconductive Topological Surface States in a Topological Superconductor

The investigation of the several potential uses of 2D topological superconductors, including quantum computing, spintronics, and quantum sensing, was the result of these investigations. Eventually, many scientists have discovered a way to induce superconducting pairing of the Dirac fermions in a two-dimensional system thereby opening up numerous possibilities for future technologies. This approach relied on the Bi2Se3/Nb heterostructure that allowed the helically pairing of the Dirac fermions and the newly found pairing of the Dirac fermions can lead to various practical applications of topological superconductors.[6] This material's two-dimensional characteristics make it ideal for quantum computing because they enable the creation of qubits that can be controlled more effectively than is feasible with traditional three-dimensional materials. Furthermore, the material is highly desirable for spintronics and quantum sensing due to the topological feature of the Dirac fermions' helical pairing, which may be used to produce numerous spin states.[7]

The image illustrates how proximity coupling with the topological superconductor as shown in figure 7 introduces superconductivity to the topological surface states of the topological insulator:

The upper layer (blue grid): a superconducting topological surface band that develops on the surface of a topological insulator when it is connected with a superconductor is shown in the upper layer (blue grid).

Lower layer (red volume): depicts a topological superconductor, which injects Cooper pairs by the proximity effect.

Green STM Tip: could be scanning tunneling microscopy (STM) or spin-polarized measurements that locally probe the superconducting surface state that is induced.

Spins & Arrows: depict spin-momentum locking and paired Majorana states at the interface that are topical for topological superconductivity.

Dashed Lines & Colored Markers: show how the topological surface states match the superconducting state underneath in terms of both space and energy.

Such a configuration is generally employed to image or electrically detect Majorana zero modes (MZMs) or induced p-wave superconductivity on the TI surface.

RESULTS AND DISCUSSION

Doped Topological Superconductors

These are materials that are composed of two distinct types of materials, one of which is a superconductor and the other a topological insulator. The physical characteristics of the resultant compound can be altered by doping both components, potentially resulting in the creation of a novel material with unique qualities that can be used in various fields.

Cu atoms are introduced between Bi2Se3 quintuple layers due to the van der Waals gaps in the Bi2Se3 structure. The intercalated Cu effectively acts as a link that binds the quintuple layers, thus enabling the phenomenon of superconductivity. The gaps between the van der Waals layers also offer room for other intercalants, i.e., Au, Pb, and Ag, all of which have been found to induce superconductivity. In other words, the superconducting characteristics of Bi2Se3 can be tuned by different doping agents. Consequently, the changed charge in the material is due to an interaction between the Bi2Se3 and the Cu dopants. The formation of Cooper pairs in the material as a result of this shift in the charge distribution is what makes Bi2Se3 superconducting.These Cooper pairs can move in the material without any resistance, hence, the superconductivity is the result.[4] Such superconductivity is also there in the case of materials like NbSe2 and YBa2Cu3O7. The effect has been heavily researched, and the studies have resulted in the discovery of the materials that can become superconductors at extremely low temperatures. For this reason, they can be employed in devices such as MRI machines and high-speed maglev trains.[5]

SrxBi2Se3

Strong chemical bonds are formed when strontium (Sr) is intercalated in Bi2Se3, making the substance more stable. One of the topological insulators is SrxBi2Se3, which has a conducting surface but an insulating bulk. Strong electron-electron interactions and topological insulator surface states in SrxBi2Se3 are thought to be responsible for the material's superconductivity. Thus, it turns out that SrxBi2Se3 is the one and only material making the diabolical duo of insulator and superconductor, which has enthralled scientists all over the globe.[6] A conduction band that is simultaneously conducting and insulating is believed to arise as a result of electron-electron interactions and the surface states of the topological insulator. It can be explained that this is why a superconductor becomes doable at much lower temperatures and higher superconductivity of 91 % volume, which is more than that of CuxBi2Se3 has been achieved so that it has become very attractive to scientists.[7] Because of this, the performance of the material is considerably enhanced, thereby making it perfect for, say, photovoltaic and electronic applications. In addition, the bond between Sr and Bi2Se3 can also be 'tuned' depending on Sr concentration, thus, allowing more control over the material's intrinsic properties as shown in figure 8. Tin-based superconductors (Sn1-xSbx, SnAs & Sn4As3).




Figure 8. XRD pattern shows some satellite peaks of SnSb systems

Quantum Sensing Advantages for Oil & Gas Monitoring

A deep dive into cutting-edge material-science research over the past few years reveals how crucial it is for nanoscale structures to be stable if the oil and gas industry wants to benefit from quantum technologies.[19] The papers mentioned show that a small change in the size of a nanoparticle can affect the XRD peak shapes and make the diffraction patterns broader to a large extent, thus pointing to the industrial materials being very sensitive to the small changes in the crystal structure. In case oil and chemical processing apparatuses, which need nanoparticle-based catalysts, corrosion-resistant coatings, or oxide interfaces, are exposed to such structural changes, they may face significant consequences in their performance. The revelations from this research constitute the ground reason why quantum-grade sensors and instruments have to be made resistant to structural drift, temperature impact, and particle size–related disorder. When reviewing quantum sensing platforms, one can hardly find any comparison between them and traditional methods in terms of methane detection, pipeline

integrity monitoring, or early-stage equipment failure identification. Thanks to their magnetic and electromagnetic sensitivity, quantum sensors, e.g., NV-center devices and superconducting magnetometers, are capable of detecting the lowest levels of a signal that cannot be reached by classical systems. It has been reported that these sensors[20] operate with high fidelity in noisy industrial environments, which is a necessary condition for their use in the field of refinery units and midstream pipeline corridors. As per the summary of conventional methane gas detection methods,[21] quantum-enhanced detectors can lead to leak identification at an earlier stage and also facilitate continuous, high-resolution mapping of emissions.

Table 1. Quantum Sensing Advantages for Oil & Gas Monitoring

Parameter / Need	Classical Technologies	Quantum-Enhanced Technologies
Methane leak detection threshold	Moderate; often ppm-level	Ultra-sensitive; potential sub-ppm capability
Resistance to noise & harsh environments	Often limited by interference	High robustness due to quantum coherence properties
Spatial resolution	Meter-scale or larger	Centimeter- to sub-centimeter mapping possible
Response time	Seconds to minutes	Near real-time, rapid signal acquisition
Suitability for continuous monitoring	Limited by drift & calibration	Strong potential for persistent, stable measurements

Industrial Challenges and Quantum Technology Opportunities




Figure 9. Conceptual Integration of Quantum Technologies in Oil-Gas Systems

Studies on large-scale methane emissions make it clear that there is a strong need for advanced detection capabilities. Methane release patterns are frequently changing and have a high spatial and temporal variability - in such conditions many traditional monitoring devices are inefficient. The implementation of quantum sensing in these processes might give operators the opportunity to detect sudden, low-intensity leak events that have not yet become large. [22] High-resolution measurement combined with real-time analytics can potentially lead to less environmental impact and help energy companies comply with stricter global emissions standards. Significantly, these discoveries illustrate that the difficulties in monitoring within the oil and gas industry are perfectly compatible with the capabilities of quantum measurement technologies.

Quantum computational tools can do more than just sensing; they have a promising potential to be able to accelerate reservoir simulation, chemical-reaction modeling, and optimization problems across the energy sector. The paper states that quantum algorithms[23] can solve high-dimensional models more efficiently than classical solvers, especially in systems that involve nonlinear coupling and uncertainty. Incorporating them in digital-twin frameworks for drilling, production forecasting, and refinery optimization, quantum simulations can possibly aid in lessening the computational delays and enhancing the predictive accuracy.[24]

As a result, the interplay of material science, quantum sensing, and quantum computation points to a future industrial ecosystem where quantum technologies function alongside classical methods, thus providing a tiered, high-precision approach to monitoring, modelling, and decision-making in oil, gas, and chemical operations.

Table 2. Industrial Challenges and Quantum Technology Opportunities

Industrial Challenge	Limitation of Current Methods	Potential Quantum Contribution	Linked Reference
Methane emissions monitoring	Missing intermittent leaks	Real-time, high-resolution mapping	Collins et al.[23]
Pipeline integrity assessment	Low sensitivity to micro-defects	Quantum magnetometry for early crack detection	Crawford et al.[21]
Catalyst performance drift	Size-dependent variation in nanoparticles	Quantum-precision materials characterization	Vorontsov et al.[20]
Reservoir simulation	Computational bottlenecks	Quantum-assisted subsurface modeling	Paudel et al.[22]
Detection of corrosive gas formation	Slow response, low precision	Quantum spectral sensors with faster detection	Aldhafeeri et al.[24]

CONCLUSIONS

A unique environment where both superconductivity and topological order occur simultaneously is provided by topological superconductors (TSCs), which represent a groundbreaking convergence of condensed matter physics and quantum information theory. By the apparition of Majorana fermions and other non-standard quasiparticles, TSCs offer a very stable road to fault-tolerant quantum computing. Moreover, their inherent capability to shield themselves from local disturbances makes them highly attractive for quantum technologies of the future. Nevertheless, probing the microscopic aspects that lead to unconventional pairing and phenomenon of symmetry-breaking is still a big puzzle to scientists.

In the very recent past, many experimental studies have singled out a variety of substances—namely Cu-doped Bi_2Se_3, Sr_2RuO_4, UTe_2, and transition metal pnictides—that demonstrate unconventional superconducting properties. Hence, these revelations have paved the route

for scientists to grasp the mechanisms by which spin–orbit coupling, symmetry principles, and electronic correlations interplay to give rise to topological superconductivity. However, the real challenge of making intrinsic TSCs in a lab is that it is only feasible under very stringent low-temperature and high-purity conditions. To unveil novel TSCs, it is therefore a continued challenge to perfect material synthesis methods like molecular beam epitaxy and hybrid heterostructure fabrication.

Besides that, the creation of hybrid and artificial TSCs has been instrumental in providing new possibilities for device-level integration. For instance, the combinations of semiconductor–superconductor nanowire systems, 2D heterostructures, and doped topological insulators are good demonstrations of how proximity-induced superconductivity may be controlled to achieve specific topological characteristics. In turn, these developments offer the quantum world a prospect of easily scalable architectures with applications in quantum computing, quantum sensing, and spintronics. Unfortunately, the issues related to reproducibility, stability, and fabrication on a large scale are still very difficult to overcome, thus delaying the inevitable practical deployment of the technologies.

Furthermore, researchers from a variety of disciplines, including condensed matter physics, materials science, and quantum engineering, must work together to develop topological superconductors in the future. Forthcoming program may entail production of high-temperature TSCs, in-depth investigation of non-centrosymmetric and moiré systems, as well as invention of highly accurate instruments capable of detecting Majorana zero modes. Once these breakthroughs get their fruits, the path will be cleared not only for the subsequent quantum devices but also, by the change of the entire modern electronics world, to a new revolution in technology to follow, with quantum technologies being super-efficient and thus computing and energy being the domains of the next technology revolution.

BIBLIOGRAPHIC REFERENCES

1. Chiu CK, Teo JC, Schnyder AP, Ryu S. Classification of topological quantum matter with symmetries. Rev Mod Phys. 2016;88(3):035005.

2. Abud M, Sartori G. The geometry of spontaneous symmetry breaking. Ann Phys. 1983;150(2):307-72.

3. Haritha N, Student PG. Introduction to Some Properties of General Topology. BIBLIOGRAPHY BIBLIOGRAPHY.

4. Marar T. A ludic journey into geometric topology. Springer International Publishing; 2022.

5. Adhikari MR. Basic algebraic topology and its applications. Springer India; 2016.

6. Alspaugh D. Majorana Quasiparticles in Topological Material Interfaces. Louisiana State University and Agricultural & Mechanical College; 2021.

7. Cyrot M. Ginzburg-Landau theory for superconductors. Rep Prog Phys. 1973;36(2):103.

8. Li Z. Free Fermion Systems: Topological Classification and Real-Space Invariants. University of Pittsburgh; 2020.

9. Tokura Y, Yasuda K, Tsukazaki A. Magnetic topological insulators. Nat Rev Phys. 2019;1(2):126-43.

10. Mandal M, Drucker NC, Siriviboon P, Nguyen T, Boonkird A, Lamichhane TN, et al. Topological superconductors from a materials perspective. Chem Mater. 2023;35(16):6184-200.

11. Luo H, Yu P, Li G, Yan K. Topological quantum materials for energy conversion and storage. Nat Rev Phys. 2022;4(9):611-24.

12. Bagchi M. Surface characterization of topological superconductor materials using scanning tunneling microscopy and spectroscopy. Universität zu **Köln; 2023.**

13. Niu Q, Thouless DJ, Wu YS. Quantized Hall conductance as a topological invariant. Phys Rev B. 1985;31(6):3372.

14. Elliott SR, Franz M. Colloquium: Majorana fermions in nuclear, particle, and solid-state physics. Rev Mod Phys. 2015;87(1):137-63.

15. Mihelic FM. Magnetic vector potential manipulation of Majorana fermions in DNA quantum logic. In: Quantum Information Science, Sensing, and Computation XIII. SPIE; 2021. p. 27-40.

16. Gruber CS, Abdel-Hafiez M. Exploring Superconductivity: The Interplay of Electronic Orders in Topological Quantum Materials. arXiv preprint arXiv:2405.17036. 2024.

17. Leijnse M, Flensberg K. Introduction to topological superconductivity and Majorana fermions. Semicond Sci Technol. 2012;27(12):124003.

18. Crogman HT, Dang T, Erenso D. Topologically Protected Quantum Teleportation via Majorana Zero Modes: A Perspective on Scalability and Decoherence Immunity. Quantum Rep. 2025;7(3):42.

19. Kou SP. Realization of topological quantum computation with planar codes. Phys Rev A. 2009;80(5):052317.

20. Vorontsov AV, Tsybulya SV. Influence of nanoparticles size on XRD patterns for small monodisperse nanoparticles of Cu0 and TiO2 anatase. Ind Eng Chem Res. 2018;57(7):2526-36.

21. Crawford SE, Shugayev RA, Paudel HP, Lu P, Ohodnicki PR, Gentry R, Duan Y. Quantum sensing for energy applications: review and perspective. Adv Quantum Technol. 2021;4(12):2100049.

22. Paudel HP, Syamlal M, Crawford SE, Lee YL, Shugayev RA, Lu P, Ohodnicki PR, Mollot D, Duan Y. Quantum computing and simulations for energy applications: review and perspective. ACS Eng Au. 2022;2(3):151–96.

23. Collins W, Orbach R, Bailey M, Biraud S, Coddington I, DiCarlo D, Peischl J, Radhakrishnan A, Schimel D, et al. Monitoring methane emissions from oil and gas operations. PRX Energy. 2022;1(1):017001.

24. Aldhafeeri T, Al-Qahtani H, Al-Ghamdi S. A review of methane gas detection sensors. Inventions. 2020;5(3):28. https://doi.org/10.3390/inventions5030028

FINANCING

None.

CONFLICT OF INTEREST

None.

AUTHORSHIP CONTRIBUTION

Conceptualization: S. Dhamotharan, M. Muthukrishnan, M. Bharath, P. Mathankumar, N. Ragavan, P. Prabhu, Shafi Danyalov.

Data curation: S. Dhamotharan, M. Muthukrishnan, M. Bharath, P. Mathankumar, N. Ragavan, P. Prabhu, Shafi Danyalov.

Formal analysis: S. Dhamotharan, M. Muthukrishnan, M. Bharath, P. Mathankumar, N. Ragavan, P. Prabhu, Shafi Danyalov.

Drafting - original draft: S. Dhamotharan, M. Muthukrishnan, M. Bharath, P. Mathankumar, N. Ragavan, P. Prabhu, Shafi Danyalov.

Writing - proofreading and editing: S. Dhamotharan, M. Muthukrishnan, M. Bharath, P. Mathankumar, N. Ragavan, P. Prabhu, Shafi Danyalov.